

APPLICATION UNDER UNITED STATES PATENT LAWS

Atty. Dkt. No. PW 279062
(M#)

Invention: NUCLEOTIDE SEQUENCES ENCODING A SENSOR KINASE, *citA*, FROM
CORYNEBACTERIUM GLUTAMICUM

Inventor (s): Bettina MÖCKEL
Mike FARWICK
Thomas HERMANN
Achim MARX
Walter PFEFFERLE



For correspondence Address



00909

Pillsbury Winthrop LLP

This is a:

- ☐ Provisional Application
- ☐ Regular Utility Application
- ☐ Continuing Application
 - ☒ The contents of the parent are incorporated by reference
- ☐ PCT National Phase Application
- ☐ Design Application
- ☐ Reissue Application
- ☐ Plant Application
- ☒ Substitute Specification
 - Sub. Spec Filed March 6, 2003
 - in App. No. 09/804,060
- ☐ Marked up Specification re
 - Sub. Spec. filed _____
 - In App. No. / _____

SPECIFICATION



**Nucleotide Sequences Encoding a Sensor Kinase,
citA, from Corynebacterium Glutamicum**

The invention provides nucleotide sequences from coryneform bacteria which code for the citA gene and a process for the fermentative preparation of amino acids, in particular L-lysine, by attenuation of the citA gene. The citA gene codes for the sensor kinase Cit A of a two-component system.

Background

10 L-Amino acids, in particular L-lysine, are used in human medicine and in the pharmaceuticals industry, in the foodstuffs industry and very particularly in animal nutrition.

It is known that amino acids are prepared by fermentation from strains of coryneform bacteria, in particular Corynebacterium glutamicum. Because of their great importance, work is constantly being undertaken to improve the preparation processes. Improvements to the process can relate to fermentation measures, such as, for example, stirring and supply of oxygen, or the composition of the nutrient media, such as, for example, the sugar concentration during the fermentation, or the working up to the product form by, for example, ion exchange chromatography, or the intrinsic output properties of the microorganism itself.

Methods of mutagenesis, selection and mutant selection are used to improve the output properties of these microorganisms. Strains which are resistant to antimetabolites or are auxotrophic for metabolites of regulatory importance and which produce amino acids are obtained in this manner.

Methods of the recombinant DNA technique have also been employed for some years for improving the strain of *Corynebacterium* strains which produce L-amino acids.

Object of the Invention

- 5 The inventors had the object of providing new measures for improved fermentative preparation of amino acids, in particular L-lysine.

Brief Description of the Invention

- 10 If L-lysine or lysine are mentioned in the following, this also means the salts, such as e.g., lysine monohydrochloride or lysine sulfate.

The invention provides an isolated polynucleotide from coryneform bacteria, comprising a polynucleotide sequence which codes for the *citA* gene chosen from the group
15 consisting of

- a) polynucleotide which is identical to the extent of at least 70% to a polynucleotide which codes for a polypeptide which comprises the amino acid sequence of SEQ ID No. 2,
- 20 b) polynucleotide which codes for a polypeptide which comprises an amino acid sequence which is identical to the extent of at least 70% to the amino acid sequence of SEQ ID No. 2,
- c) polynucleotide which is complementary to the
25 polynucleotides of a) or b), and
- d) polynucleotide comprising at least 15 successive nucleotides of the polynucleotide sequence of a), b) or c),

the polypeptide preferably having the activity of sensor
30 kinase CitA.

The invention also provides the abovementioned polynucleotide, this preferably being a DNA which is capable of replication, comprising:

- (i) the nucleotide sequence shown in SEQ ID No. 1 or
- 5 (ii) at least one sequence which corresponds to sequence (i) within the range of the degeneration of the genetic code, or
- (iii) at least one sequence which hybridizes with the sequences complementary to sequences (i) or (ii),
10 and optionally
- (iv) sense mutations of neutral function in (i).

The invention also provides:

a polynucleotide, in particular DNA, which is capable of replication and comprises the nucleotide sequence as
15 shown in SEQ ID No. 1;

a polynucleotide which codes for a polypeptide which comprises the amino acid sequence as shown in SEQ ID No. 2;

a vector containing parts of the polynucleotide according
20 to the invention, but at least 15 successive nucleotides of the sequence claimed

and coryneform bacteria in which the citA gene is attenuated, in particular by an insertion or deletion.

The invention also provides polynucleotides which
25 substantially comprise a polynucleotide sequence, which are obtainable by screening by means of hybridization of a corresponding gene library of a coryneform bacterium, which comprises the complete gene or parts thereof, with a probe which comprises the sequence of the polynucleotide
30 according to the invention or a fragment thereof, and isolation of the polynucleotide sequence mentioned.

Polynucleotides comprising the sequences according to the invention are suitable as hybridization probes for RNA, cDNA and DNA, in order to isolate, in the full length, nucleic acids or polynucleotides or genes which code for
5 Cita protein or to isolate those nucleic acids or polynucleotides or genes which have a high similarity with the sequence of the cita gene.

Polynucleotides comprising the sequences according to the invention are furthermore suitable as primers with the aid
10 of which DNA of genes which code for the Cita protein can be prepared by the polymerase chain reaction (PCR).

Such oligonucleotides which serve as probes or primers comprise at least 30, preferably at least 20, very particularly preferably at least 15 successive nucleotides.
15 Oligonucleotides which have a length of at least 40 or 50 nucleotides are also suitable.

Drawing
Detailed Description of the ~~Invention~~

The following figure is attached:

Figure 1: Map of the plasmid pCR2.1citAint.

20 The abbreviations and designations used have the following meaning.

KmR:	Kanamycin resistance gene
EcoRI:	Cleavage site of the restriction enzyme EcoRI
HindIII:	Cleavage site of the restriction enzyme HindIII
BamHI:	Cleavage site of the restriction enzyme BamHI
citAint:	Internal fragment of the cita gene

KK
5/23/03

ColE1: Replication origin of the plasmid ColE1

Detailed Description of the Invention

"Isolated" means separated out of its natural environment.

"Polynucleotide" in general relates to polyribonucleotides and polydeoxyribonucleotides, it being possible for these
5 to be non-modified RNA or DNA or modified RNA or DNA.

The polynucleotides according to the invention include a polynucleotide according to SEQ ID No. 1 or a fragment prepared therefrom and also those which are at least 70%, preferably at least 80% and in particular at least 90% to
10 95% identical to the polynucleotide according to SEQ ID No. 1 or a fragment prepared therefrom.

"Polypeptides" are understood as meaning peptides or proteins which comprise two or more amino acids bonded via peptide bonds.

15 The polypeptides according to the invention include a polypeptide according to SEQ ID No. 2, in particular those with the biological activity of the CitA protein and also those which are at least 70%, preferably at least 80% and in particular at least 90% to 95% identical to the
20 polypeptide according to SEQ ID No. 2 and have the activity mentioned.

The invention moreover relates to a process for the fermentative preparation of amino acids, in particular L-lysine, using coryneform bacteria which in particular
25 already produce amino acids and in which the nucleotide sequences which code for the citA gene are attenuated, in particular eliminated or expressed at a low level.

The term "attenuation" in this connection describes the reduction or elimination of the intracellular activity of
30 one or more enzymes (proteins) in a microorganism which are coded by the corresponding DNA, for example by using a weak

KK
5/23/03

promoter or using a gene or allele which codes for a corresponding enzyme with a low activity or inactivates the corresponding gene or enzyme (protein), and optionally combining these measures.

- 5 The microorganisms which the present invention provides can prepare amino acids, in particular L-lysine, from glucose, sucrose, lactose, fructose, maltose, molasses, starch, cellulose or from glycerol and ethanol. They can be representatives of coryneform bacteria, in particular of
10 the genus *Corynebacterium*. Of the genus *Corynebacterium*, there may be mentioned in particular the species *Corynebacterium glutamicum*, which is known among experts for its ability to produce L-amino acids.

Suitable strains of the genus *Corynebacterium*, in
15 particular of the species *Corynebacterium glutamicum* (*C. glutamicum*), are in particular the known wild-type strains

- Corynebacterium glutamicum* ATCC13032
Corynebacterium acetoglutamicum ATCC15806
Corynebacterium acetoacidophilum ATCC13870
20 *Corynebacterium melassecola* ATCC17965
Corynebacterium thermoaminogenes FERM BP-1539
Brevibacterium flavum ATCC14067
Brevibacterium lactofermentum ATCC13869 and
Brevibacterium divaricatum ATCC14020

- 25 or L-amino acid-producing mutants or strains prepared therefrom, such as, for example, the L-lysine-producing strains

- Corynebacterium glutamicum* FERM-P 1709
Brevibacterium flavum FERM-P 1708
30 *Brevibacterium lactofermentum* FERM-P 1712
Corynebacterium glutamicum FERM-P 6463
Corynebacterium glutamicum FERM-P 6464
Corynebacterium glutamicum DM58-1
Corynebacterium glutamicum DG52-5

Corynebacterium glutamicum DSM 5715 and
Corynebacterium glutamicum DSM 12866

The inventors have succeeded in isolating the new *citA* gene of *C. glutamicum* which codes for the CitA protein and which is a sensor kinase of a two-component system.

To isolate the *citA* gene or also other genes of *C. glutamicum*, a gene library of this microorganism is first set up in *Escherichia coli* (*E. coli*). The setting up of gene libraries is described in generally known textbooks and handbooks. The textbook by Winnacker: *Genes and Clones, An Introduction to Genetic Engineering* (Verlag Chemie, Weinheim, Germany, 1990), or the handbook by Sambrook et al.: *Molecular Cloning, A Laboratory Manual* (Cold Spring Harbor Laboratory Press, 1989) may be mentioned as an example. A well-known gene library is that of the *E. coli* K-12 strain W3110 set up in λ -vectors by Kohara et al. (*Cell* 50, 495-508 (1987)). Bathe et al. (*Molecular and General Genetics*, 252:255-265, 1996) describe a gene library of *C. glutamicum* ATCC13032, which was set up with the aid of the cosmid vector SuperCos I (Wahl et al., 1987, *Proceedings of the National Academy of Sciences USA*, 84:2160-2164) in the *E. coli* K-12 strain NM554 (Raleigh et al., 1988, *Nucleic Acids Research* 16:1563-1575). Börmann et al. (*Molecular Microbiology* 6(3), 317-326 (1992)) in turn describe a gene library of *C. glutamicum* ATCC13032 using the cosmid pH79 (Hohn and Collins, 1980, *Gene* 11, 291-298).

To prepare a gene library of *C. glutamicum* in *E. coli* it is also possible to use plasmids such as pBR322 (Bolivar, 1979, *Life Sciences*, 25, 807-818) or pUC9 (Vieira et al., 1982, *Gene*, 19:259-268). Suitable hosts are, in particular, those *E. coli* strains which are restriction- and recombination-defective, such as, for example, the strain DH5 α (Jeffrey H. Miller: "A Short Course in Bacterial Genetics, A Laboratory Manual and Handbook for

Escherichia coli and Related Bacteria," Cold Spring Harbor Laboratory Press, 1992).

The long DNA fragments cloned with the aid of cosmids or other λ -vectors can then be subcloned in turn into the
5 usual vectors suitable for DNA sequencing.

Methods of DNA sequencing are described, inter alia, by Sanger et al. (Proceedings of the National Academy of Sciences of the United States of America USA, 74:5463-5467, 1977).

10 The resulting DNA sequences can then be investigated with known algorithms or sequence analysis programs, such as e.g., that of Staden (Nucleic Acids Research 14, 217-232(1986)), that of Marck (Nucleic Acids Research 16, 1829-1836 (1988)) or the GCG program of Butler (Methods of
15 Biochemical Analysis 39, 74-97 (1998)).

The new DNA sequence of *C. glutamicum* which codes for the *citA* gene and which, as SEQ ID No. 1, is a constituent of the present invention has been found in this way. The amino acid sequence of the corresponding protein has
20 furthermore been derived from the present DNA sequence by the methods described above. The resulting amino acid sequence of the *citA* gene product is shown in SEQ ID No. 2.

Coding DNA sequences which result from SEQ ID No. 1 by the degeneracy of the genetic code are also a constituent of
25 the invention. In the same way, DNA sequences which hybridize with SEQ ID No. 1 or parts of SEQ ID No. 1 are a constituent of the invention. Conservative amino acid exchanges, such as e.g., exchange of glycine for alanine or of aspartic acid for glutamic acid in proteins, are
30 furthermore known among experts as "sense mutations" which do not lead to a fundamental change in the activity of the protein, i.e., are of neutral function. It is furthermore known that changes on the N and/or C terminus of a protein cannot substantially impair or can even stabilize the

function thereof. Information in this context can be found by the expert, inter alia, in Ben-Bassat et al. (Journal of Bacteriology 169:751-757 (1987)), in O'Regan et al. (Gene 77:237-251 (1989)), in Sahin-Toth et al. (Protein Sciences 5 3:240-247 (1994)), in Hochuli et al. (Bio/Technology 6:1321-1325 (1988)) and in known textbooks of genetics and molecular biology. Amino acid sequences which result in a corresponding manner from SEQ ID No. 2 are also a constituent of the invention.

10 Finally, DNA sequences which are prepared by the polymerase chain reaction (PCR) using primers which result from SEQ ID No. 1 are a constituent of the invention. Such oligonucleotides typically have a length of at least 15 nucleotides.

15 Instructions for identifying DNA sequences by means of hybridization can be found by the expert, inter alia, in the handbook "The DIG System Users Guide for Filter Hybridization" from Boehringer Mannheim GmbH (Mannheim, Germany, 1993) and in Liebl et al. (International Journal
20 of Systematic Bacteriology 41: 255-260 (1991)). The hybridization takes place under stringent conditions, that is to say only hybrids in which the probe and target sequence, i.e., the polynucleotides treated with the probe, are at least 70% identical are formed. It is known that
25 the stringency of the hybridization, including the washing steps, is influenced or determined by varying the buffer composition, the temperature and the salt concentration. The hybridization reaction is preferably carried out under a relatively low stringency compared with the washing steps
30 (Hybaid Hybridisation Guide, Hybaid Limited, Teddington, UK, 1996). A 5x SSC buffer at a temperature of approx. 50 - 68°C, for example, can be employed for the hybridization reaction. Probes can also hybridize here with polynucleotides which are less than 70% identical to the
35 sequence of the probe. Such hybrids are less stable and are removed by washing under stringent conditions. This

can be achieved, for example, by lowering the salt concentration to 2x SSC and optionally subsequently 0.5x SSC (The DIG System User's Guide for Filter Hybridisation, Boehringer Mannheim, Mannheim, Germany, 1995) a temperature of approx. 50 - 68°C being established. Polynucleotide fragments which are, for example, at least 70% or at least 80% or at least 90% to 95% identical to the sequence of the probe employed can be isolated by increasing the hybridization temperature stepwise from 50 to 68°C in steps of approx. 1 - 2°C. Further instructions on hybridization are obtainable on the market in the form of so-called kits (e.g., DIG Easy Hyb from Roche Diagnostics GmbH, Mannheim, Germany, Catalogue No. 1603558).

Instructions for amplification of DNA sequences with the aid of the polymerase chain reaction (PCR) can be found by the expert, inter alia, in the handbook by Gait: Oligonucleotides: A Practical Approach (IRL Press, Oxford, UK, 1984) and in Newton and Graham: PCR (Spektrum Akademischer Verlag, Heidelberg, Germany, 1994).

In the work on the present invention, it has been found that coryneform bacteria produce amino acids, in particular L-lysine, in an improved manner after attenuation of the *citA* gene.

To achieve an attenuation, either the expression of the *citA* gene or the catalytic properties of the enzyme protein can be reduced or eliminated. The two measures can optionally be combined.

The reduction in gene expression can take place by suitable culturing or by genetic modification (mutation) of the signal structures of gene expression. Signal structures of gene expression are, for example, repressor genes, activator genes, operators, promoters, attenuators, ribosome binding sites, the start codon and terminators.

The expert can find information on this e.g., in the patent application WO 96/15246, in Boyd and Murphy (Journal of

Bacteriology 170: 5949 (1988)), in Voskuil and Chambliss (Nucleic Acids Research 26: 3548 (1998), in Jensen and Hammer (Biotechnology and Bioengineering 58: 191 (1998)), in Pátek et al. (Microbiology 142: 1297 (1996)), Vasicova
5 et al. (Journal of Bacteriology 181: 6188 (1999)) and in known textbooks of genetics and molecular biology, such as e.g., the textbook by Knippers ("Molecular Genetics," 6th edition, Georg Thieme Verlag, Stuttgart, Germany, 1995) or that by Winnacker ("Genes and Clones," VCH
10 Verlagsgesellschaft, Weinheim, Germany, 1990).

Mutations which lead to a change or reduction in the catalytic properties of enzyme proteins are known from the prior art; examples which may be mentioned are the works by Qiu and Goodman (Journal of Biological Chemistry 272: 8611-
15 8617 (1997)), Sugimoto et al. (Bioscience Biotechnology and Biochemistry 61: 1760-1762 (1997)) and Möckel ("Threonine dehydratase from Corynebacterium glutamicum: Cancelling the allosteric regulation and structure of the enzyme," Reports from the Jülich Research Centre, Jül-2906,
20 ISSN09442952, Jülich, Germany, 1994). Summarizing descriptions can be found in known textbooks of genetics and molecular biology, such as e.g., that by Hagemann ("General Genetics," Gustav Fischer Verlag, Stuttgart, 1986).

25 Possible mutations are transitions, transversions, insertions and deletions. Depending on the effect of the amino acid exchange on the enzyme activity, missense mutations or nonsense mutations are referred to. Insertions or deletions of at least one base pair (bp) in a
30 gene lead to frame shift mutations, as a consequence of which incorrect amino acids are incorporated or translation is interrupted prematurely. Deletions of several codons typically lead to a complete loss of the enzyme activity. Instructions on generation of such mutations are prior art
35 and can be found in known textbooks of genetics and molecular biology, such as e.g., the textbook by Knippers

("Molecular Genetics," 6th edition, Georg Thieme Verlag, Stuttgart, Germany, 1995), that by Winnacker ("Genes and Clones," VCH Verlagsgesellschaft, Weinheim, Germany, 1990) or that by Hagemann ("General Genetics", Gustav Fischer Verlag, Stuttgart, 1986).

A common method of mutating genes of *C. glutamicum* is the method of gene disruption and gene replacement described by Schwarzer and Pühler (Bio/Technology 9, 84-87 (1991)).

In the method of gene disruption a central part of the coding region of the gene of interest is cloned in a plasmid vector which can replicate in a host (typically *E. coli*), but not in *C. glutamicum*. Possible vectors are, for example, pSUP301 (Simon et al., Bio/Technology 1, 784-791 (1983)), pK18mob or pK19mob (Schäfer et al., Gene 145, 69-73 (1994)), pK18mobsacB or pK19mobsacB (Jäger et al., Journal of Bacteriology 174: 5462-65 (1992)), pGEM-T (Promega corporation, Madison, WI, USA), pCR2.1-TOPO (Shuman (1994). Journal of Biological Chemistry 269:32678-84; US Patent 5,487,993), pCR®Blunt (Invitrogen, Groningen, Holland; Bernard et al., Journal of Molecular Biology, 234: 534-541 (1993)) or pEM1 (Schrumpf et al, 1991, Journal of Bacteriology 173:4510-4516). The plasmid vector which contains the central part of the coding region of the gene is then transferred into the desired strain of *C. glutamicum* by conjugation or transformation. The method of conjugation is described, for example, by Schäfer et al. (Applied and Environmental Microbiology 60, 756-759 (1994)). Methods for transformation are described, for example, by Thierbach et al. (Applied Microbiology and Biotechnology 29, 356-362 (1988)), Dunican and Shivan (Bio/Technology 7, 1067-1070 (1989)) and Tauch et al. (FEMS Microbiological Letters 123, 343-347 (1994)). After homologous recombination by means of a "cross-over" event, the coding region of the gene in question is interrupted by the vector sequence and two incomplete alleles are obtained, one lacking the 3' end and one lacking the 5'

end. This method has been used, for example, by Fitzpatrick et al. (Applied Microbiology and Biotechnology 42, 575-580 (1994)) to eliminate the *recA* gene of *C. glutamicum*.

5 In the method of gene replacement, a mutation, such as e.g., a deletion, insertion or base exchange, is established in vitro in the gene of interest. The allele prepared is in turn cloned in a vector which is not replicative for *C. glutamicum* and this is then transferred
10 into the desired host of *C. glutamicum* by transformation or conjugation. After homologous recombination by means of a first "cross-over" event which effects integration and a suitable second "cross-over" event which effects excision in the target gene or in the target sequence, the
15 incorporation of the mutation or of the allele is achieved. This method was used, for example, by Peters-Wendisch et al. (Microbiology 144, 915 - 927 (1998)) to eliminate the *pyc* gene of *C. glutamicum* by a deletion.

A deletion, insertion or a base exchange can be
20 incorporated into the *citA* gene in this manner.

In addition, it may be advantageous for the production of L-amino acids, in particular L-lysine, to enhance, in particular to over-express, one or more enzymes of the particular biosynthesis pathway, of glycolysis, of
25 anaplerosis, of the pentose phosphate cycle or of amino acid export and optionally regulatory proteins, in addition to attenuation of the *citA* gene.

The term "enhancement" in this connection describes the increase in the intracellular activity of one or more
30 enzymes (proteins) in a microorganism which are coded by the corresponding DNA, for example by increasing the number of copies of the gene or genes, using a potent promoter or using a gene or allele which codes for a corresponding enzyme (protein) having a high activity, and optionally
35 combining these measures.

Thus, for example, for the preparation of L-lysine, at the same time one or more of the genes chosen from the group consisting of

- the dapA gene which codes for dihydrodipicolinate
5 synthase (EP-B 0 197 335),
 - the gap gene which codes for glyceraldehyde 3-phosphate dehydrogenase (Eikmanns (1992), Journal of Bacteriology 174:6076-6086),
 - the zwf gene which codes for glucose 6-phosphate
10 dehydrogenase (JP-A-09224661),
 - the pyc gene which codes for pyruvate carboxylase (DE-A-198 31 609),
 - the lysE gene which codes for lysine export (DE-A-195 48 222),
 - 15 • the lysC gene which codes for a feed back resistant aspartate kinase (EP-B-0387527; EP-A-0699759; WO 00/63388), or
 - the zwal gene which codes for the Zwal protein (DE: 199 59 328.0, DSM 13115)
- 20 can be enhanced, in particular over-expressed.

It may furthermore be advantageous for the production of amino acids, in particular L-lysine, in addition to the attenuation of the citA gene, at the same time for one or more of the genes chosen from the group consisting of

- 25 • the pck gene which codes for phosphoenol pyruvate carboxykinase (DE 199 50 409.1, DSM 13047),
- the pgi gene which codes for glucose 6-phosphate isomerase (US 09/396,478, DSM 12969),

- the poxB gene which codes for pyruvate oxidase (DE:1995 1975.7, DSM 13114),
- the zwa2 gene which codes for the Zwa2 protein (DE: 199 59,327.2, DSM 13113)

5 to be attenuated.

In addition to attenuation of the citA gene it may furthermore be advantageous, for the production of amino acids, in particular L-lysine, to eliminate undesirable side reactions, (Nakayama: "Breeding of Amino Acid
10 Producing Microorganisms," in: Overproduction of Microbial Products, Krumphanzl, Sikyta, Vanek (eds.), Academic Press, London, UK, 1982).

The invention also provides the microorganisms prepared according to the invention, and these can be cultured
15 continuously or discontinuously in the batch process (batch culture) or in the fed batch (feed process) or repeated fed batch process (repetitive feed process) for the purpose of production of L-amino acids, in particular L-lysine. A summary of known culture methods is described in the
20 textbook by Chmiel (Bioprocess Technology 1. Introduction to Bioprocess Technology (Gustav Fischer Verlag, Stuttgart, 1991)) or in the textbook by Storhas (Bioreactors and Peripheral Equipment (Vieweg Verlag, Braunschweig/Wiesbaden, 1994)).

25 The culture medium to be used must meet the requirements of the particular strains in a suitable manner. Descriptions of culture media for various microorganisms are contained in the handbook "Manual of Methods for General Bacteriology" of the American Society for Bacteriology
30 (Washington D.C., USA, 1981). Sugars and carbohydrates, such as e.g., glucose, sucrose, lactose, fructose, maltose, molasses, starch and cellulose, oils and fats, such as, for example, soya oil, sunflower oil, groundnut oil and coconut fat, fatty acids, such as, for example, palmitic acid,

stearic acid and linoleic acid, alcohols, such as, for example, glycerol and ethanol, and organic acids, such as, for example, acetic acid, can be used as the source of carbon. These substances can be used individually or as a
5 mixture.

Organic nitrogen-containing compounds, such as peptones, yeast extract, meat extract, malt extract, corn steep liquor, soya bean flour and urea, or inorganic compounds, such as ammonium sulfate, ammonium chloride, ammonium
10 phosphate, ammonium carbonate and ammonium nitrate, can be used as the source of nitrogen. The sources of nitrogen can be used individually or as a mixture.

Phosphoric acid, potassium dihydrogen phosphate or dipotassium hydrogen phosphate or the corresponding sodium-
15 containing salts can be used as the source of phosphorus. The culture medium must furthermore comprise salts of metals, such as, for example, magnesium sulfate or iron sulfate, which are necessary for growth. Finally, essential growth substances, such as amino acids and
20 vitamins, can be employed in addition to the abovementioned substances. Suitable precursors can moreover be added to the culture medium. The starting substances mentioned can be added to the culture in the form of a single batch, or can be fed in during the culture in a suitable manner.

25 Basic compounds, such as sodium hydroxide, potassium hydroxide, ammonia or aqueous ammonia, or acid compounds, such as phosphoric acid or sulfuric acid, can be employed in a suitable manner to control the pH of the culture. Antifoams, such as, for example, fatty acid polyglycol
30 esters, can be employed to control the development of foam. Suitable substances having a selective action, such as, for example, antibiotics, can be added to the medium to maintain the stability of plasmids. To maintain aerobic conditions, oxygen or oxygen-containing gas mixtures, such
35 as, for example, air, are introduced into the culture. The temperature of the culture is usually 20°C to 45°C, and

preferably 25°C to 40°C. Culturing is continued until a maximum of the desired product has formed. This target is usually reached within 10 hours to 160 hours.

Methods for the determination of L-amino acids are known
5 from the prior art. The analysis can thus be carried out, for example, as described by Spackman et al. (Analytical Chemistry, 30, (1958), 1190) by anion exchange chromatography with subsequent ninhydrin derivatization, or
it can be carried out by reversed phase HPLC, for example
10 as described by Lindroth et al. (Analytical Chemistry (1979) 51: 1167-1174).

The invention furthermore relates to a process for the fermentative preparation of an amino acid chosen from the group consisting of L-asparagine, L-threonine, L-serine, L-
15 glutamate, L-glycine, L-alanine, L-cysteine, L-valine, L-methionine, L-isoleucine, L-leucine, L-tyrosine, L-phenylalanine, L-histidine, L-lysine, L-tryptophan and L-arginine, in particular L-lysine, using coryneform bacteria which in particular already produce one or more of the
20 amino acids mentioned.

The following microorganism was deposited on 23.01.2001 as a pure culture at the German Collection of Microorganisms and Cell Cultures, Braunschweig, Germany in accordance with the Budapest Treaty:

- 25 • Escherichia coli strain Top10/pCR2.1citAint as DSM 13998.

The present invention is explained in more detail in the following with the aid of embodiment examples.

The isolation of plasmid DNA from Escherichia coli and all
30 techniques of restriction, Klenow and alkaline phosphatase treatment were carried out by the method of Sambrook et al. (Molecular Cloning. A Laboratory Manual, 1989, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, USA).

Methods for transformation of *Escherichia coli* are also described in this handbook.

The composition of the usual nutrient media, such as LB or TY medium, can also be found in the handbook by Sambrook et al.

ExamplesExample 1

Preparation of a genomic cosmid gene library from *C. glutamicum* ATCC 13032

- 5 Chromosomal DNA from *C. glutamicum* ATCC 13032 was isolated as described by Tauch et al. (1995, Plasmid 33:168-179) and partly cleaved with the restriction enzyme Sau3AI (Amersham Pharmacia, Freiburg, Germany, Product Description Sau3AI, Code no. 27-0913-02). The DNA fragments were
- 10 dephosphorylated with shrimp alkaline phosphatase (Roche Molecular Biochemicals, Mannheim, Germany, Product Description SAP, Code no. 1758250). The DNA of the cosmid vector SuperCos1 (Wahl et al. (1987), Proceedings of the National Academy of Sciences, USA 84:2160-2164), obtained
- 15 from Stratagene (La Jolla, USA, Product Description SuperCos1 Cosmid Vektor Kit, Code no. 251301) was cleaved with the restriction enzyme XbaI (Amersham Pharmacia, Freiburg, Germany, Product Description XbaI, Code no. 27-0948-02) and likewise dephosphorylated with shrimp alkaline
- 20 phosphatase.

The cosmid DNA was then cleaved with the restriction enzyme BamHI (Amersham Pharmacia, Freiburg, Germany, Product Description BamHI, Code no. 27-0868-04). The cosmid DNA treated in this manner was mixed with the treated ATCC13032

25 DNA and the batch was treated with T4 DNA ligase (Amersham Pharmacia, Freiburg, Germany, Product Description T4-DNA-Ligase, Code no. 27-0870-04). The ligation mixture was then packed in phages with the aid of Gigapack II XL Packing Extract (Stratagene, La Jolla, USA, Product

30 Description Gigapack II XL Packing Extract, Code no. 200217).

For infection of the *E. coli* strain NM554 (Raleigh et al. 1988, Nucleic Acid Res. 16:1563-1575) the cells were taken up in 10 mM MgSO₄ and mixed with an aliquot of the phage

suspension. The infection and titering of the cosmid library were carried out as described by Sambrook et al. (1989, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor), the cells being plated out on LB agar (Lennox, 5 1955, Virology, 1:190) + 100 µg/ml ampicillin. After incubation overnight at 37°C, recombinant individual clones were selected.

Example 2

Isolation and sequencing of the *citA* gene

10 The cosmid DNA of an individual colony was isolated with the Qiaprep Spin Miniprep Kit (Product No. 27106, Qiagen, Hilden, Germany) in accordance with the manufacturer's instructions and partly cleaved with the restriction enzyme Sau3AI (Amersham Pharmacia, Freiburg, Germany, Product 15 Description Sau3AI, Product No. 27-0913-02). The DNA fragments were dephosphorylated with shrimp alkaline phosphatase (Roche Molecular Biochemicals, Mannheim, Germany, Product Description SAP, Product No. 1758250). After separation by gel electrophoresis, the cosmid 20 fragments in the size range of 1500 to 2000 bp were isolated with the QiaExII Gel Extraction Kit (Product No. 20021, Qiagen, Hilden, Germany).

The DNA of the sequencing vector pZero-1, obtained from Invitrogen (Groningen, The Netherlands, Product Description 25 Zero Background Cloning Kit, Product No. K2500-01) was cleaved with the restriction enzyme BamHI (Amersham Pharmacia, Freiburg, Germany, Product Description BamHI, Product No. 27-0868-04). The ligation of the cosmid fragments in the sequencing vector pZero-1 was carried out 30 as described by Sambrook et al. (1989, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor), the DNA mixture being incubated overnight with T4 ligase (Pharmacia Biotech, Freiburg, Germany). This ligation mixture was then electroporated (Tauch et al. 1994, FEMS Microbiol 35 Letters, 123:343-7) into the *E. coli* strain DH5αMCR (Grant,

1990, Proceedings of the National Academy of Sciences, U.S.A., 87:4645-4649) and plated out on LB agar (Lennox, 1955, Virology, 1:190) with 50 µg/ml zeocin.

The plasmid preparation of the recombinant clones was
5 carried out with Biorobot 9600 (Product No. 900200, Qiagen, Hilden, Germany). The sequencing was carried out by the dideoxy chain termination method of Sanger et al. (1977, Proceedings of the National Academies of Sciences, U.S.A., 74:5463-5467) with modifications according to Zimmermann et
10 al. (1990, Nucleic Acids Research, 18:1067). The "RR dRhodamin Terminator Cycle Sequencing Kit" from PE Applied Biosystems (Product No. 403044, Weiterstadt, Germany) was used. The separation by gel electrophoresis and analysis of the sequencing reaction were carried out in a
15 "Rotiphoresis NF Acrylamide/Bisacrylamide" Gel (29:1) (Product No. A124.1, Roth, Karlsruhe, Germany) with the "ABI Prism 377" sequencer from PE Applied Biosystems (Weiterstadt, Germany).

The raw sequence data obtained were then processed using
20 the Staden program package (1986, Nucleic Acids Research, 14:217-231) version 97-0. The individual sequences of the pZerol derivatives were assembled to a continuous contig. The computer-assisted coding region analysis was prepared with the XNIP program (Staden, 1986, Nucleic Acids
25 Research, 14:217-231). Further analyses were carried out with the "BLAST search program" (Altschul et al., 1997, Nucleic Acids Research, 25:3389-3402) against the non-redundant databank of the "National Center for Biotechnology Information" (NCBI, Bethesda, MD, USA).

30 The resulting nucleotide sequence is shown in SEQ ID No. 1. Analysis of the nucleotide sequence showed an open reading frame of 1653 bp, which was called the *citA* gene. The *citA* gene codes for a polypeptide of 551 amino acids.

Example 3

Preparation of an integration vector for integration mutagenesis of the *citA* gene

From the strain ATCC 13032, chromosomal DNA was isolated by the method of Eikmanns et al. (Microbiology 140: 1817 - 1828 (1994)). On the basis of the sequence of the *citA* gene known for *C. glutamicum* from example 2, the following oligonucleotides were chosen for the polymerase chain reaction (see SEQ ID No. 4 and SEQ ID No. 5):

10 *citA*-int1:

5' TTC CAG TCG GTG AGG TCA GT 3'

citA-int2:

5' GTA CGA TCG CGG ATG GTT AC 3'

The primers shown were synthesized by MWG Biotech (Ebersberg, Germany) and the PCR reaction was carried out by the standard PCR method of Innis et al. (PCR protocols. A guide to methods and applications, 1990, Academic Press) with the Taq-polymerase from Boehringer Mannheim (Germany, Product Description Taq DNA polymerase, Product No. 1 146 165). With the aid of the polymerase chain reaction, the primers allow amplification of an internal fragment of the *citA* gene 480 bp in size. The product amplified in this way was tested electrophoretically in a 0.8% agarose gel.

The amplified DNA fragment (see SEQ ID No. 3) was ligated with the TOPO TA Cloning Kit from Invitrogen Corporation (Carlsbad, CA, USA; Catalogue Number K4500-01) in the vector pCR2.1-TOPO (Mead et al. (1991) Bio/Technology 9:657-663).

The *E. coli* strain TOP10 was then electroporated with the ligation batch (Hanahan, In: DNA cloning. A practical approach, Vol. I, IRL-Press, Oxford, Washington DC, USA, 1985). Selection for plasmid-carrying cells was made by plating out the transformation batch on LB agar (Sambrook et al., Molecular cloning: a laboratory manual. 2nd Ed.

Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989), which had been supplemented with 50 mg/l kanamycin. Plasmid DNA was isolated from a transformant with the aid of the QIAprep Spin Miniprep Kit from Qiagen and checked by restriction with the restriction enzyme EcoRI and subsequent agarose gel electrophoresis (0.8%). The plasmid was called pCR2.1citAint and is shown in figure 1.

The following microorganism was deposited as a pure culture on 23.01.2001 at the German Collection of Microorganisms and Cell Cultures, Braunschweig, Germany in accordance with the Budapest Treaty:

- *Escherichia coli* Top10/pCR2.1citAint as DSM 13998

Example 4

Integration mutagenesis of the *citA* gene in the strains DSM 5715 and FERM-BP 1763

The vector pCR2.1citAint mentioned in example 3 was electroporated by the electroporation method of Tauch et al. (FEMS Microbiological Letters, 123:343-347 (1994)) into the strains *Corynebacterium glutamicum* DSM 5715 and *Brevibacterium lactofermentum* FERM-BP 1763. The strain DSM 5715 is an AEC-resistant lysine producer (EP-B-0435132), and the strain FERM-BP 1763 is a mycophenolic acid-resistant valine producer (US-A-5,188,948). The vector pCR2.1citAint cannot replicate independently in DSM5715 and FERM-BP 1763 and is retained in the cell only if it has integrated into the chromosome of DSM 5715 or FERM-BP 1763. Selection of clones with pCR2.1citAint integrated into the chromosome was carried out by plating out the electroporation batch on LB agar (Sambrook et al., Molecular cloning: a laboratory manual. 2nd Ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.), which had been supplemented with 15 mg/l kanamycin.

For detection of the integration, the citAint fragment was labelled with the Dig hybridization kit from Boehringer by the method of "The DIG System Users Guide for Filter Hybridization" of Boehringer Mannheim GmbH (Mannheim, Germany, 1993). Chromosomal DNA of in each case a potential integrant was isolated by the method of Eikmanns et al. (Microbiology 140: 1817 - 1828 (1994)) and in each case cleaved with the restriction enzymes EcoRI, BamHI and HindIII. The fragments formed were separated by means of agarose gel electrophoresis and hybridized at 68°C with the Dig hybridization kit from Boehringer. The plasmid pCR2.1citAint mentioned in example 3 had been inserted into the chromosome of DSM5715 and into the chromosome of FERM-BP 1763 within the chromosomal citA gene. The strains were called DSM5715::pCR2.1citAint and FERM-BP 1763::pCR2.1citAint.

Example 5

Preparation of lysine

The *C. glutamicum* strain DSM5715::pCR2.1citAint obtained in example 4 was cultured in a nutrient medium suitable for the production of lysine and the lysine content in the culture supernatant was determined.

For this, the strain was first incubated on an agar plate with the corresponding antibiotic (brain-heart agar with kanamycin (25 mg/l) for 24 hours at 33°C. Starting from this agar plate culture, a preculture was seeded (10 ml medium in a 100 ml conical flask). The complete medium CgIII was used as the medium for the preculture.

Medium Cg III

NaCl	2.5 g/l
Bacto-Peptide	10 g/l
Bacto-Yeast extract	10 g/l
Glucose (autoclaved separately)	2% (w/v)

The pH is brought to pH 7.4

Kanamycin (25 mg/l) was added to this. The preculture was incubated for 16 hours at 33°C at 240 rpm on a shaking
5 machine. A main culture was seeded from this preculture such that the initial OD (660 nm) of the main culture is 0.1 OD. Medium MM was used for the main culture.

Medium MM

CSL (corn steep liquor)	5 g/l
MOPS (morpholinopropanesulfonic acid)	20 g/l
Glucose (autoclaved separately)	50g/l
Salts:	
(NH ₄) ₂ SO ₄	25 g/l
KH ₂ PO ₄	0.1 g/l
MgSO ₄ * 7 H ₂ O	1.0 g/l
CaCl ₂ * 2 H ₂ O	10 mg/l
FeSO ₄ * 7 H ₂ O	10 mg/l
MnSO ₄ * H ₂ O	5.0 mg/l
Biotin (sterile-filtered)	0.3 mg/l

Thiamine * HCl (sterile-filtered)	0.2 mg/l
Leucine (sterile-filtered)	0.1 g/l
CaCO ₃	25 g/l

The CSL, MOPS and the salt solution were brought to pH 7 with aqueous ammonia and autoclaved. The sterile substrate and vitamin solutions were then added, and the CaCO₃ 5 autoclaved in the dry state was added.

Culturing was carried out in a 10 ml volume in a 100 ml conical flask with baffles. Kanamycin (25 mg/l) was added. Culturing was carried out at 33°C and 80% atmospheric humidity.

10 After 72 hours, the OD was determined at a measurement wavelength of 660 nm with a Biomek 1000 (Beckmann Instruments GmbH, Munich). The amount of lysine formed was determined with an amino acid analyzer from Eppendorf-BioTronik (Hamburg, Germany) by ion exchange chromatography 15 and post-column derivatization with ninhydrin detection.

The result of the experiment is shown in table 1.

Table 1

Strain	OD(660)	Lysine HCl g/l
DSM5715	7.5	13.3
DSM5715::pCR2.1citAint	7.4	14.4

Example 6

Preparation of valine

The *B. lactofermentum* strain FERM-BP 1763::pCR2.1citAint obtained in example 4 was cultured in a nutrient medium
5 suitable for the production of valine and the valine content in the culture supernatant was determined.

For this, the strain was first incubated on an agar plate with the corresponding antibiotic (brain-heart agar with kanamycin (25 mg/l) for 24 hours at 33°C. Starting from
10 this agar plate culture, a preculture was seeded (10 ml medium in a 100 ml conical flask). The complete medium CgIII was used as the medium for the preculture. Kanamycin (25 mg/l) was added to this. The preculture was incubated for 16 hours at 33°C at 240 rpm on a shaking machine. A
15 main culture was seeded from this preculture such that the initial OD (660nm) of the main culture was 0.1 OD. Medium MM was used for the main culture.

Medium MM

CSL (corn steep liquor)	5 g/l
MOPS (morpholinopropanesulfonic acid)	20 g/l
Glucose (autoclaved separately)	50 g/l
Salts:	
(NH ₄) ₂ SO ₄	25 g/l
KH ₂ PO ₄	0.1 g/l
MgSO ₄ * 7 H ₂ O	1.0 g/l
CaCl ₂ * 2 H ₂ O	10 mg/l
FeSO ₄ * 7 H ₂ O	10 mg/l

MnSO ₄ * H ₂ O	5.0 mg/l
Isoleucine (sterile-filtered)	0.1 g/l
Methionine (sterile-filtered)	0.1 g/l
Thiamine * HCl (sterile-filtered)	0.2 mg/l
Leucine (sterile-filtered)	0.1 g/l
CaCO ₃	25 g/l

The CSL (corn steep liquor), MOPS
(morpholinopropanesulfonic acid) and the salt solution were
brought to pH 7 with aqueous ammonia and autoclaved. The
5 sterile substrate and vitamin solutions were then added,
and the CaCO₃ autoclaved in the dry state was added.

Culturing was carried out in a 10 ml volume in a 100 ml
conical flask with baffles. Kanamycin (25 mg/l) was added.
Culturing was carried out at 33°C and 80% atmospheric
10 humidity.

After 48 hours, the OD was determined at a measurement
wavelength of 660 nm with a Biomek 1000 (Beckmann
Instruments GmbH, Munich). The amount of valine formed was
determined with an amino acid analyzer from Eppendorf-
15 BioTronik (Hamburg, Germany) by ion exchange chromatography
and post-column derivatization with ninhydrin detection.

The result of the experiment is shown in table 2.

Table 2

Strain	OD(660)	Valine HCl g/l
FERM-BP 1763	12.1	7.5
FERM-BP 1763::pCR2.1citAint	13.5	10.8